

# If only I could tell ...

Citation for published version (APA):

Buskermolen, W. M. (2017). If only I could tell ...: measuring predictors for challenging behaviour in people with both intellectual disability and hearing impairment. [Doctoral Thesis, Maastricht University]. Maastricht University. <https://doi.org/10.26481/dis.20170504wb>

**Document status and date:**

Published: 04/05/2017

**DOI:**

[10.26481/dis.20170504wb](https://doi.org/10.26481/dis.20170504wb)

**Document Version:**

Publisher's PDF, also known as Version of record

**Document license:**

Unspecified

**Please check the document version of this publication:**

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)

**Take down policy**

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.

## Summary

Our study with 3,5-diiodo-L-thyronine (T2), a thyroid hormone that was earlier regarded functionally inactive, has revealed its effects in the two-day tail suspension test. In this model, a bolus injection of T2 at two different doses, to mice of two different strains, resulted in a decrease of depressive-like behaviour, thus, suggesting the antidepressant-like effect of T2 (**Chapter 2**). This effect of T2 was not known before; it suggested clinical use of T2 under conditions that require faster than with other thyroid hormones therapeutic effect. However, no effects of T2 on depressive-like behaviour were found in a classic two-day forced swim test (**Chapter 3**). Thus, our results have demonstrated a limited sensitivity of the forced swim test that is in line with available literature, and suggested a need for its modification so the accuracy of the test could be increased.

In this context, we modified commonly used protocol of the forced swim test, where an additional session on Day 5 has followed the initial exposure on Days 1 and 2 in its classical variant (**Chapter 4**). We hypothesized that additional delayed session on Day 5 can model increased contextual learning and consolidation of environmental adversities related to the testing. Increased brain expression of GSK3 $\beta$ , a hallmark of a distress and depression in humans was observed in mice after modified, but not classic sessions of the swim test (**Chapter 4**). Moreover, increased scores of depressive-like behaviour (floating behaviour) during additional delayed testing positively correlated with the levels of GSK3 $\beta$  mRNA in the prefrontal cortex of experimental groups (**Chapter 4**). No such correlations were found in the hippocampal formation, while GSK3 $\beta$  expression was increased there as well. Elevated activities of GSK3 $\beta$  during repeated but not single exposure to swimming were further demonstrated on a protein level. The concentration of 9-Ser-phosphorylated (inactive) form of pS9-GSK3 $\beta$  and its ratio to total GSK3 $\beta$  were significantly decreased during repeated testing (**Chapter 4**).

Our experiments have identified the role of timing and exposure to a context in the development of behavioural and molecular changes during the modified swim test. A “premature” exposure to swimming on Day 3 instead

of Day 5 did not result in expression of GSK3 $\beta$  (**Chapter 4**). On other hand, a replacement of a swim session with an exposure to the context of testing on Day 2 and Day 5 or the omission of intermediate swimming or context exposure on these days affected floating behaviour and brain GSK3 $\beta$  mRNA in the modified swim test (**Chapter 4**). Together, our experiments suggested the critical role of GSK3 $\beta$  in the “consolidation phase” of the development depressive-like features associated with enhanced conditioning to adverse context.

Two-week delivery of low dose of imipramine (7.5 mg/kg/day), thiamine and benfotiamine (200 mg/kg/day) has precluded an increase of floating and over-expression of in mice during the last session of the modified forced swim test (**Chapter 5**). Given that all these compounds were shown to decrease GSK3 $\beta$  activities, these data further suggest the role of this molecular factor in enhanced contextual conditioning associated with depressive-like state. In addition, these data have identified thiamine and its precursor as antidepressant agents and suggested their usefulness in treatment of depression in clinic (**Chapter 5**).